Clinical Risk Factors for Severe Clostridium difficile-associated **Disease**

Timothy J. Henrich, Douglas Krakower, Asaf Bitton, and Deborah S. Yokoe

CME ACTIVITY

Medscape, LLC is pleased to provide online continuing medical education (CME) for this journal article, allowing clinicians the opportunity to earn CME credit. This activity has been planned and implemented in accordance with the Essential Areas and policies of the Accreditation Council for Continuing Medical Education through the joint sponsorship of Medscape, LLC and Emerging Infectious Diseases. Medscape, LLC is accredited by the Accreditation Council for Continuing Medical Education (ACCME) to provide CME for physicians. Medscape, LLC designates this educational activity for a maximum of 0.75 AMA PRA Category 1 CreditsTM. Physicians should only claim credit commensurate with the extent of their participation in the activity. All other clinicians completing this activity will be issued a certi?cate of participation. To participate in this journal CME activity: (1) review the learning objectives and author disclosures; (2) study the education content; (3) take the post-test and/ or complete the evaluation at http://www.medscape.com/cme/eid; (4) view/print certi?cate.

Learning Objectives

Upon completion of this activity, participants will be able to:

- Identify the criteria used to de?ne severe Clostridium difficile-associated disease (CDAD) in the current study
- Specify the prevalence of severe CDAD in the current study
- Identify the clinical risk factors for severe CDAD
- List the laboratory risk factors for severe CDAD

Lynne Stockton, Technical Writer-Editor, Emerging Infectious Diseases. Disclosure: Lynne Stockton has disclosed no relevant financial

Charles P. Vega, MD, Associate Professor; Residency Director, Department of Family Medicine, University of California, Irvine, California, USA. Disclosure: Charles P. Vega, MD, has disclosed that he has served as an advisor or consultant to Novartis, Inc.

Disclosures: Timothy J. Henrich, MD; Douglas Krakower, MD; Asaf Bitton, MD; and Deborah S. Yokoe, MD, MPH, have disclosed no relevant financial relationships.

Identifying patients who are at high risk for severe Clostridium difficile-associated disease (CDAD) early in the course of their infection may help clinicians improve outcomes. Therefore, we compared clinical features associated with severe versus nonsevere CDAD by retrospectively reviewing records of hospitalized patients whose fecal assays were positive for C. difficile toxin. Of 336 patients, 12.2% had severe disease and 10.1% died from all causes. Regression modeling showed the following to be significantly associated with severe CDAD (p≤0.05): age >70 years (odds ratio [OR] 3.35), maximum leukocyte count >20,000 cells/mL (OR 2.77), minimum albumin level <2.5 g/dL (OR 3.44), maximum creatinine level >2 mg/dL (OR 2.47), small bowel obstruction or ileus (OR 3.06), and computed tomography scan showing colorectal inflammation (OR 13.54). These clinical and laboratory markers for severe disease

Author affiliations: Brigham and Women's Hospital, Boston, Massachusetts, USA (T.J. Henrich, D. Krakower, A. Bitton, D.S. Yokoe); Massachusetts General Hospital, Boston (D. Krakower); and Harvard Medical School, Boston (Deborah S. Yokoe)

DOI: 10.3201/eid1503.080312

may be useful for identifying patients at risk for serious outcomes or death.

The incidence and severity of *Clostridium difficile*—asso-L ciated disease (CDAD) is increasing in North America (1-3) and Europe (4,5). During the past 10 years in the United States, prevalence, case-fatality rates, total attributable mortality rates, and colectomy rates for persons with CDAD have markedly increased (6). Acquisition of C. difficile and the development of severe CDAD is associated primarily with healthcare, although community-acquired severe disease among persons previously thought to be at low risk for infection have been reported (5,7,8). Several mechanisms for increased disease severity have been proposed, including emergence of speci? c strains with genetic polymorphisms that encode higher levels of bacterial toxins A and B and the production of a binary toxin (3,9,10). The Centers for Disease Control and Prevention has reported outbreaks of CDAD associated with the new BI/ NAP1 strain in 40 of 50 US states, although the association

Emerging Infectious Diseases • www.cdc.gov/eid • Vol. 15, No. 3, March 2009

415

Earning CME Credit

To obtain credit, you should ?rst read the journal article. After reading the article, you should be able to answer the following, related, multiple-choice questions. To complete the questions and earn continuing medical education (CME) credit, please go to http://www.medscape.com/cme/eid. Credit cannot be obtained for tests completed on paper, although you may use the worksheet below to keep a record of your answers. You must be a registered user on Medscape.com. If you are not registered on Medscape.com, please click on the New Users: Free Registration link on the left hand side of the website to register. Only one answer is correct for each question. Once you successfully answer all post-test questions you will be able to view and/or print your certi?cate. For questions regarding the content of this activity, contact the accredited provider, CME@medscape.net. For technical assistance, contact CME@webmd.net. American Medical Association's Physician's Recognition Award (AMA PRA) credits are accepted in the US as evidence of participation in CME activities. For further information on this award, please refer to http://www.ama-assn.org/ama/pub/category/2922. html. The AMA has determined that physicians not licensed in the US who participate in this CME activity are eligible for AMA PRA Category 1 CreditsTM. Through agreements that the AMA has made with agencies in some countries, AMA PRA credit is acceptable as evidence of participation in CME activities. If you are not licensed in the US and want to obtain an AMA PRA CME credit, please complete the questions online, print the certi? cate and present it to your national medical association.

Article Title

Clinical Risk Factors for Severe Clostridium difficileassociated Disease

CME Questions

- 1. All of the following were criteria for severe Clostridium difficile-associated disease (CDAD) in the current study, except:
- A. One or more intensive care unit admissions in which C. difficile was a major contributor
- B. Prolonged symptoms past 14 days requiring intravenous fluid replacement
- C. Colectomy or other surgery directly attributed to C. difficile
- D. Intestinal perforation in the setting of *C. difficile* infection
- 2. What was the prevalence of severe CDAD among all of the cases of CDAD in the current study?
- A. <1%
- B. 12%
- C. 29% D. 44%

- 3. Which of the following patient factors was most associated with an increased risk for severe CDAD on multivariate analysis of the current study?
- Age >70 years
- B. Chemotherapy use
- C. Antimicrobial use D. Previous hospital stay
- 4. All of the following laboratory factors were predictive of an increased risk for CDAD in the current study, except:
- A. White blood cell count >20,000 cells/mL
- B. Serum albumin <2.5 g/dL
- C. Creatinine >2 mg/dL
- D. Alanine aminotransferase >40 U/L

Activity Evaluation

 The activity supported the Strongly Disagree 	learning objectives.			Strongly Agree
1	2	3	4	5
. The material was organize	d clearly for learning	to occur.		
Strongly Disagree				Strongly Agree
1	2	3	4	5
. The content learned from	this activity will impa	ct my practice.		
Strongly Disagree				Strongly Agree
1	2	3	4	5
. The activity was presented	d objectively and free	of commercial bias.		
Strongly Disagree	-			Strongly Agree
1	2	3	4	5

Emerging Infectious Diseases • www.cdc.gov/eid • Vol. 15, No. 3, March 2009